



"Pack year" smoking histories: what about patients who use loose tobacco?

D M Wood, M G Mould, S B Y Ong and E H Baker

Tob. Control 2005;14:141-142
doi:10.1136/tc.2004.009977

Updated information and services can be found at:
<http://tc.bmjournals.com/cgi/content/full/14/2/141>

These include:

References

This article cites 10 articles, 3 of which can be accessed free at:
<http://tc.bmjournals.com/cgi/content/full/14/2/141#BIBL>

Rapid responses

You can respond to this article at:
<http://tc.bmjournals.com/cgi/eletter-submit/14/2/141>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

[Smoking](#) (1002 articles)
[Tobacco use](#) (160 articles)
[Health effects of tobacco use](#) (109 articles)

Notes

To order reprints of this article go to:
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *Tobacco Control* go to:
<http://www.bmjournals.com/subscriptions/>

BRIEF REPORT

“Pack year” smoking histories: what about patients who use loose tobacco?

D M Wood, M G Mould, S B Y Ong, E H Baker

Tobacco Control 2005;14:141–142. doi: 10.1136/tc.2004.009977

Life time smoking of ready made cigarettes can be readily quantified as “pack years” smoked, but methods for measuring loose tobacco use are less well established. In this study the frequency of loose tobacco use by 247 hospital in-patients was determined; 64% were current or ex-smokers, 41.3% of whom (25.9% of participants) had smoked loose tobacco. A formula was developed for converting loose tobacco use to pack years smoked, based on the weight of tobacco in ready made cigarettes; 12.5 g or half an ounce of loose tobacco was approximately equivalent to one packet of 20 cigarettes. Using a questionnaire it was found that hospital physicians of all grades were able to convert smoking histories of ready made cigarettes, but not loose tobacco, into number of “pack years” smoked.

Lifetime smoking exposure is quantified in “pack years”, where one “pack year” is 20 cigarettes smoked/day for one year.¹ Quantification of pack years smoked is important in clinical care where degree of tobacco exposure is closely correlated to risk of disease. Smokers suffer an irreversible forced expiratory volume in one second (FEV₁) loss of 4.4–10.4 ml per pack year smoked.^{2–3} There is a strong dose–response relation between the number of pack years smoked and the risk,⁴ severity,⁵ and mortality⁶ of chronic obstructive pulmonary disease and the risk of lung cancer.⁷ Pack year smoking histories are important for quantification of tobacco exposure in research for direct study, to define patient groups or to determine confounding effects of smoking on outcomes from other exposures.

Although the number of ready made cigarettes smoked can easily be quantified in pack years, there is no well established method for measuring loose tobacco use. In this study we

determined whether there was a need to quantify loose tobacco exposure and developed a simple method for converting loose tobacco use into pack years.

PARTICIPANTS, METHODS, AND RESULTS

Prevalence of previous or current loose tobacco use was investigated by a two day cross sectional survey of hospital in-patients > 16 years old. Age, sex, in-patient speciality, smoking history, and lifetime loose tobacco use of participants were recorded. The relation between ready made cigarettes and loose tobacco was estimated by weight. The ability of physicians of different grades (table 1) to calculate pack years was assessed by questionnaire. Scenarios for ready made cigarettes (n = 5) and loose tobacco (n = 5) were given for conversion into total number of pack years smoked.

A total of 333 hospital in-patients were approached and 247 (74%) completed the survey. Non-participants were confused (n = 17), unwilling (n = 14), or too unwell (n = 55). Of the participants, 22.7% were current smokers, 41.3% were ex-smokers, and 36% had never smoked; 41.3% of current/ex-smokers and 25.9% of all participants had smoked loose tobacco. Males (40.5%) were more likely than females to be current/ex-smokers (males 71%, females 58.3%, p = 0.029, χ^2 test) or have smoked loose tobacco (males 62.9%, females 23.2%, p < 0.0001). Participants under medical or surgical consultants were equally likely to be current/ex-smokers (p = 0.34) or have smoked loose tobacco (p = 0.24). Ages were similar in those who had (mean (SD) 62.0 (18.6) years) or had not smoked loose tobacco (61.8 (20.4) years, p = 0.45, unpaired t test).

The mean (SD) weight of tobacco per ready made cigarette was 0.73 (0.029) (n = 20) and 20 cigarettes approximated a 12.5 g (1/2 ounce) sachet of loose tobacco. Loose tobacco users usually describe tobacco smoked in “ounces per week”, which can be converted into pack years:

Table 1 Responses of medical and surgical pre-registration house officers (PRHO), medical senior house officers (SHO), respiratory specialist registrars (SpR), and general medicine consultants to history taking questionnaire and scenario assessment. Results are based on returned questionnaires

	Medical/ surgical PRHO	Medical SHO	Respiratory SpR	Medical consultant
Number surveyed	23	11	18	17
Number of respondents (%)	18 (78.3)	11 (100)	16 (88.9)	10 (55.6)
Converts ready made cigarette use into “pack years” in clinical practice (% of respondents)	13 (72.2)	4 (36.3)	14 (87.5)	3 (30)
Converts loose tobacco use into “pack years” in clinical practice (% of respondents)	0 (0)	0 (0)	3 (18.8)	2 (20)
Scenario score (maximum 5) for conversion of ready made cigarettes into pack years (mean (SD) of respondents)	4.2 (1.1)	3.9 (1.9)	4.6 (0.6)	3.2 (2.3)
Scenario score (maximum 5) for conversion of loose tobacco use into pack years (mean (SD) of respondents)	0 (0)	0 (0)	0.2 (0.4)	0 (0)

ounces per week \times 2/7 \times number of years smoked = pack years

Fifty five of 69 (79.7%) physicians approached completed the questionnaire (table 1). Physicians were able to and commonly did convert ready made cigarettes into pack years (score $4.1 \pm 1.4/5$). Physicians were unable to quantify loose tobacco use (score $0 \pm 0.2/5$).

DISCUSSION

We have shown that many smokers use loose tobacco, and that clinicians are unable to quantify loose tobacco use. We have therefore developed a simple formula to allow quantification of loose tobacco and ready made cigarette consumption in the same "units" for estimation of total smoking load. This formula has application both in clinical practice and research.

The formula has limitations as it does not allow for different smoking habits. Loose tobacco is more commonly smoked without a filter than ready made cigarettes, and smoking habits are determined by tar and nicotine content of the tobacco.^{8,9} Despite this we feel our formula is useful because of the strong dose-response relation between total tobacco consumption and ill health.^{2,7} Other variables such as inhalation and use of filters have not consistently been shown to influence clinical outcomes.^{7,10}

Authors' affiliations

D M Wood, Pharmacology and Clinical Pharmacology, St George's Hospital Medical School, London, UK
M G Mould, S B Y Ong, St George's Hospital Medical School
E H Baker, Physiological Medicine, St George's Hospital Medical School

Correspondence to: Dr Emma Baker, Physiological Medicine, St George's Hospital Medical School, Cranmer Terrace, London SW17 0RE, UK; ebaker@sghms.ac.uk

Received 13 September 2004
 Accepted 23 November 2004

What this paper adds

There is no well established method for quantifying lifetime use of loose tobacco. We found that loose tobacco use was common in hospital in-patients, implying that measurement of this use was desirable. We therefore developed a simple weight based conversion of loose tobacco into "pack years" smoked which should be used in clinical practice and research: ounces per week \times 2/7 \times number of years smoked = pack years

REFERENCES

- 1 **Prignot J**. Quantification and chemical markers of tobacco-exposure. *Eur J Respir Dis* 1987;**70**:1-7.
- 2 **Dockery DW**, Speizer FE, Ferris BG, *et al*. Cumulative and reversible effects of lifetime smoking on simple tests of lung function in adults. *Am Rev Respir Dis* 1988;**137**:286-92.
- 3 **Prescott E**, Bjerg AM, Anderson PK, *et al*. Gender difference in smoking effects on lung function and risk of hospitalization for COPD: results from a Danish longitudinal population study. *Eur Respir J* 1997;**10**:822-7.
- 4 **Jaen Diaz JI**, de Castro Mesa C, Gontan Garcia-Salamanca MJ, *et al*. Prevalence of chronic obstructive pulmonary disease and risk factors in smokers and ex-smokers. *Arch Bronconeumol* 2003;**39**:554-8.
- 5 **Kornmann O**, Beeh KM, Beier J, *et al*. Global initiative for obstructive lung disease. Newly diagnosed chronic obstructive pulmonary disease. Clinical features and distribution of the novel stages of the global initiative for obstructive lung disease. *Respiration* 2003;**70**:67-75.
- 6 **Kanner RE**, Renzetti AD Jr, Stanish WM, *et al*. Predictors of survival in subjects with chronic airflow limitation. *Am J Med* 1983;**74**:249-55.
- 7 **Agudo A**, Ahrens W, Benhamou E, *et al*. Lung cancer and cigarette smoking in women: a multicenter case-control study in Europe. *Int J Cancer* 2000;**88**:820-7.
- 8 **Wald NJ**, Idle M, Boreham J, *et al*. Inhaling habits among smokers of different types of cigarettes. *Thorax* 1980;**35**:925-8.
- 9 **Wald NJ**, Boreham J, Bailey A. Relative intakes of tar, nicotine, and carbon monoxide from cigarettes of different yields. *Thorax* 1984;**39**:361-4.
- 10 **Beck GJ**, Doyle CA, Schachter EN. Smoking and lung function. *Am Rev Respir Dis* 1981;**123**:149-55.

The Lighter Side



A November 2004 poll in London's *Daily Mirror* shows an encouraging shift in public opinion on public place smoking bans.