



## Interventions to prevent back pain and back injury in nurses: a systematic review

Anna P Dawson, Skye N McLennan, Stefan D Schiller, Gwendolen A Jull, Paul W Hodges and Simon Stewart

*Occup. Environ. Med.* 2007;64;642-650; originally published online 23 May 2007;

doi:10.1136/oem.2006.030643

---

Updated information and services can be found at:  
<http://oem.bmj.com/cgi/content/full/64/10/642>

---

*These include:*

### References

This article cites 45 articles, 6 of which can be accessed free at:  
<http://oem.bmj.com/cgi/content/full/64/10/642#BIBL>

1 online articles that cite this article can be accessed at:  
<http://oem.bmj.com/cgi/content/full/64/10/642#otherarticles>

### Rapid responses

You can respond to this article at:  
<http://oem.bmj.com/cgi/eletter-submit/64/10/642>

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

---

### Notes

---

To order reprints of this article go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to *Occupational and Environmental Medicine* go to:  
<http://journals.bmj.com/subscriptions/>

## REVIEW

# Interventions to prevent back pain and back injury in nurses: a systematic review

Anna P Dawson, Skye N McLennan, Stefan D Schiller, Gwendolen A Jull, Paul W Hodges, Simon Stewart

*Occup Environ Med* 2007;**64**:642–650. doi: 10.1136/oem.2006.030643

A systematic literature review was undertaken to assess the effectiveness of interventions that aim to prevent back pain and back injury in nurses. Ten relevant databases were searched; these were examined and reference lists checked. Two reviewers applied selection criteria, assessed methodological quality and extracted data from trials. A qualitative synthesis of evidence was undertaken and sensitivity analyses performed. Eight randomised controlled trials and eight non-randomised controlled trials met eligibility criteria. Overall, study quality was poor, with only one trial classified as high quality. There was no strong evidence regarding the efficacy of any interventions aiming to prevent back pain and injury in nurses. The review identified moderate level evidence from multiple trials that manual handling training in isolation is not effective and multidimensional interventions are effective in preventing back pain and injury in nurses. Single trials provided moderate evidence that stress management programs do not prevent back pain and limited evidence that lumbar supports are effective in preventing back injury in nurses. There is conflicting evidence regarding the efficacy of exercise interventions and the provision of manual handling equipment and training. This review highlights the need for high quality randomised controlled studies to examine the effectiveness of interventions to prevent back pain and injury in nursing populations. Implications for future research are discussed.

as personality and the presence of psychosomatic symptoms. Work task and work organisational factors have been shown to be significant risks in individual studies, although when all trials are considered the evidence is inconsistent. Nursing qualifications are important, with nursing assistants at greater risk for back pain than registered nurses. Years in the nursing profession may also be relevant, with a growing body of evidence suggesting that younger nurses are at greatest risk.<sup>5</sup> Identification of individual physical predictors of back pain is more elusive. Prospective studies find predominantly non-significant relationships or inconsistent results.<sup>6–10</sup> However, reduced lateral bending of the spine has been identified as a risk factor in two studies.<sup>11 12</sup>

Back pain and injury have a major impact on the efficiency of the nursing workforce. Registered nurses rank seventh and nursing aides and orderlies are highest ranked across all occupations for back injuries involving days away from work in private industry.<sup>13</sup> Back injuries and resultant workers' compensation claims in nurses are expensive. In long-term care facilities in the United States, nurses' back injuries are estimated to cost over US\$6 million in indemnity and medical payments. Nurses' compensation for back injury comprises 56.4% of all indemnity costs and 55.1% of all medical costs.<sup>3</sup> In one Australian state, nurse back injury claims accounted for \$A2.39 million expenditure in one financial year.<sup>14</sup>

Surprisingly, few systematic reviews have investigated the efficacy of interventions to prevent back pain and injury in nurses. One review which included trials over a 10-year period up to 1998 provided a narrative summary but failed to score the methodological quality of trials or interpret findings in consideration of varying design flaws.<sup>15</sup> A systematic review of interventions to reduce musculoskeletal injuries associated with handling patients included many relevant trials but was not focused on nurses and accepted low-quality designs such as professional opinion.<sup>16</sup>

The primary objective of this study was to systematically review the literature to determine whether there are interventions with proven efficacy that prevent back pain and back injury in nurses. A review focusing specifically on nurses was considered necessary due to the unique nature of nursing work.

Nurses play an important role within the health care system, providing and assisting in the provision of primary, secondary and tertiary level health care. Typically, their work is physically demanding. Nurses frequently assist patients to mobilise, transfer between positions and perform other activities of daily living such as toileting and showering.

Nurses have an increased risk of back trouble. Compared with other professions, they have an increased risk of back pain<sup>1</sup> and a six times higher prevalence of back injury.<sup>2</sup> Nurses and related medical workers lead all other occupations for risk of herniated lumbar intervertebral discs requiring hospitalisation in women.<sup>3</sup> Furthermore, hospitals and nursing and residential care facilities lead all industries for workplace injury and illness.<sup>4</sup>

Numerous factors have been found to increase nurses' risk of back pain.<sup>5</sup> Physical load and work posture play a role, as do psychosocial factors such

See end of article for authors' affiliations

Correspondence to:  
Ms Anna Dawson, Division of Physiotherapy, School of Health and Rehabilitation Sciences, The University of Queensland, St Lucia, 4072, Brisbane, Australia; a.dawson1@uq.edu.au

Accepted 4 May 2007  
Published Online First  
23 May 2007

## METHODS

The methodological design of this review followed guidelines developed by the Editorial Board of the Cochrane Collaboration Back Review Group.<sup>17 18</sup> The guidelines directed a five-step process that involved a comprehensive literature search strategy, application of selection criteria to identify eligible trials, grading of methodological quality, extraction of data and synthesis of evidence. Unlike other reviews, we reviewed only prospective trials with concurrent control groups, scored the quality of trials, differentiated between high and low quality studies, and synthesised evidence on the basis of study quality. As there is frequent indiscriminate use of terminology, we chose to review all trials focussing on low back pain (LBP), back pain and back injury to comprehensively examine the efficacy of interventions that aim to prevent back trouble in nurses. In order to identify and minimise (where possible) any potential publication biases, searches were performed in a large number of databases with no language restriction.

### Search strategy for identification of studies

Relevant studies were identified using a comprehensive search strategy that included:

1. A computer aided search of relevant databases including MEDLINE (1966–October 2004), EMBASE (1993–November 2004), CINAHL (1982–October 2004), Academic Search Elite (1985–October 2004), Health Source Nursing/Academic Edition (1975–October 2004), PEDro (1929–September 2004), PsycInfo (1840–October 2004), PsycArticles (1985–October 2004), Joanna Briggs Institute Systematic Review Database for Evidence Based Nursing and Midwifery (1998–September 2004) databases and the Cochrane Central Register of Controlled Trials (1898–September 2004). Searches were undertaken without language or year restriction using medical subject headings, key words and free text words where necessary.
2. A search of PhD dissertations using the Australian Digital Thesis Program (1997–November 2004), Index to Theses (1970–November 2004) and Digital Dissertations (1861–November 2004) databases.
3. Screening reference lists of included papers for further eligible studies.

The search terms and strategies utilised in each of the databases are available from the corresponding author.

### Criteria for considering studies for this review Studies

Due to the language skills of the reviewing team, controlled trials that were randomised (RCTs) or non-randomised (NCTs) and published in English or German were included without year restriction. Both RCTs and NCTs were accepted in order to maximise the pool of available data. Studies needed to be of prospective design with comparison intervention and control groups recruited from the same setting.

### Participants

All grades of nurses, nursing aides and nursing students were included without gender or age limitation. Studies that included nurses as well as non-nursing participants were excluded if they did not analyse and report the nursing cohort separately. There was no restriction on history of back pain and back injury.

### Interventions

Interventions aimed at the prevention of LBP, back pain and/or back injury were included. Both singular and multidimensional

strategies were eligible. Laboratory testing of patient handling techniques or equipment was excluded.

### Outcome measures

To be eligible, studies had to assess LBP and/or back pain and/or back injury outcomes. Secondary outcomes of interest included functional status (eg, disability, restricted work and leisure activities) and time loss from work.

### Selection, quality assessment and analysis

#### Study selection

Three reviewers (AD, SM, SS) participated in study selection. For each database, two authors independently screened hits and applied selection criteria. SS translated German papers for independent assessment by the other two reviewers. Selection criteria had been piloted on a sample of abstracts using title, abstract and keywords. For those studies where eligibility was unclear from the title and abstract, the full text of the article was obtained and assessed. To avoid multiple publication bias, efforts were made to identify any multiple publications arising from single studies and pool data.

#### Methodological quality assessment

The methodological quality of studies was assessed by AD and SM. All grading was performed independently with any disagreements between authors resolved via a consensus meeting. Blinding reviewers to author and publication details was not possible as reviewers had been involved in screening databases and selecting studies. One of the reviewers was trained in an unrelated health discipline and was unfamiliar with the back pain literature.

The methodological quality of studies was evaluated using criteria recommended in relevant guidelines<sup>17 18</sup> and adapted for the study designs included in the review. Criteria regarding blinding were omitted as in many interventions it was not possible to blind the subject or care giver to the treatment, and blinding the outcome assessor was irrelevant as outcomes were self-reported. A criterion assessing compliance was omitted as in many cases no control treatment was applied, and compliance was rarely measured in the intervention group. Criteria were pilot tested on a related but ineligible paper. The “internal validity” and “descriptive quality” of trials were evaluated using six criteria and each given a score out of 6 (table 1). A total score out of 12 was assigned after combining these two values. Criteria were graded as “yes” (1), “no” (0) or “don’t know” (0). A grade of “don’t know” was assigned when data were unavailable (eg, weren’t collected) and/or when information provided in the paper and by authors in personal correspondence was insufficient to score the criteria. For example, an author merely stating that randomisation was performed was inadequate to score a “yes” for criterion A. Rather, description of the specific method of randomisation utilised was required. Studies were graded according to quality assessment scores. Studies scoring a minimum of 4.5/6 for internal validity with a total score of 9/12 or greater were deemed “high quality” with all remaining studies graded as “low quality”.

When papers provided insufficient information to assess criteria, authors of included studies were contacted via email or post to request further information. When contact information was not given on included papers or was out of date, current contact details were identified by searching for more recent publications of the author in MEDLINE and searching on general internet search engines. Second authors were approached if the first author of the paper could not be contacted or did not reply. Where author responses were unclear, they were asked to provide further information and clarification.

### Data extraction and analysis

Two reviewers (AD and SM) independently extracted data using a standardised and piloted form. Studies were grouped according to intervention method and then the strength of evidence was evaluated using a qualitative method of best-evidence synthesis for data analysis. This method evaluates the number, methodological quality and outcomes of studies to provide an overall evidence rating for each intervention type ( $n = 5$  levels). Quality of evidence was graded as follows:

- Strong: consistent findings among multiple high quality RCTs
- Moderate: one high quality RCT or consistent findings among multiple low quality RCTs/NCTs
- Limited: one low quality RCT or NCT
- Conflicting: inconsistent findings among multiple trials (RCTs/NCTs)
- No evidence from any trials.

Two sensitivity analyses were undertaken to investigate how sensitive the results of the review were in relation to the way it was performed. The first sensitivity analysis investigated the impact of the selected cut-off point for qualification as a high quality trial. In this analysis, the cut-off point was amended to a minimum score of 3/6 for both internal validity and descriptive criteria. The second sensitivity analysis examined whether evidence findings changed when more potentially biased trials were omitted from analysis. Here, studies scoring less than 2/6 for internal validity were excluded from the synthesis of evidence.

### Study selection

On the basis of title and/or abstract information, 51 papers were identified. Thirty one of these were deemed ineligible following review of the full paper. Reasons for exclusion included lack of a control group, inadequate control, lack of between-group comparison, non-nursing subjects, laboratory testing and failing to report back pain or injury symptoms separately or at all. One study was ineligible on the basis of publication language.<sup>19</sup>

Twenty papers reporting nine RCTs and 11 NCTs met selection criteria. Studies with incomplete randomisation of participants were classified as NCTs.<sup>20</sup> Multiple publications from individual trials occurred in two instances, with four

papers<sup>21–24</sup> reporting a Canadian NCT and two papers<sup>25–26</sup> presenting data from a Swedish RCT. Data given in multiple publications were pooled and are referred to by the first paper published. After pooling, eight RCTs and eight NCTs were included. Figure 1 shows the number of trials at each stage of the review process.

### Study quality

Quality appraisal results are presented in table 2. Further information to enable grading of quality was required for all trials, and all authors (first or second) were contactable and responded. In two instances, authors were not able to provide information, and in some cases authors did not answer all questions. Lack of necessary information resulted in criteria being assigned a grade of “don’t know”. There was initial disagreement between reviewers on 56 (22%) of 255 scored items. This was due to reading or interpretation errors and was resolved via consensus meeting.

Mean internal validity scores were 3.1/6 (range 1.5–4.5) for RCTs and 0.6/6 (range 0–2.0) for NCTs. Mean descriptive quality scores were 3.9/6 (range 1.5–5.5) for RCTs and 3.3/6 (range 2.5–4.0) for NCTs. Of 16 trials included, only one RCT qualified as high quality according to our criteria.<sup>27</sup>

### Study characteristics

Included studies are presented in tables 3–5 according to intervention and listed under subcategories in a hierarchical order by internal validity score. Nurses, nursing aides, student nurses and at times a combination of nurse categories comprised study cohorts. In relation to outcomes, eight trials reported LBP,<sup>25 27–33</sup> one reported back pain,<sup>20</sup> two reported back injury<sup>34 35</sup> and five reported a combination of outcomes.<sup>11 21 36–38</sup>


### Analysis of intervention efficacy

#### Exercise

We identified three RCTs and two NCTs of exercise interventions that aimed to prevent LBP in nurses (table 3). Three trials reported no effect,<sup>27–29</sup> including a high quality trial of an individually designed home exercise program.<sup>27</sup> One trial of a 13-month physiotherapist-led exercise program undertaken in the workplace reported a significant reduction in LBP prevalence and intensity.<sup>30</sup> In the study by Wigaeus Hjelm and colleagues<sup>25</sup> the results were unclear: different findings

**Table 1** Methodological quality criteria

	Score
<b>Internal validity (IV) criteria</b>	
A Method of randomisation adequate (eg, random number table)?	1
B Treatment allocation concealed?	1
C Groups similar at baseline regarding age, gender, history of back pain, area of work, pregnancy, nurse grade and baseline measures?	1
D Co-interventions avoided or similar?	1
E i) Drop-out/withdrawal individuals adequately described?	0.5
ii) Drop-out rate acceptable (<20% for short-term and <30% for long-term follow-up)?	0.5
F Intention to treat method of analysis (irrespective of compliance, change in group status)?	1
IV score	6
<b>Descriptive quality (DQ) criteria</b>	
G Eligibility criteria specified?	1
H Index and control interventions explicitly described?	1
I i) Short-term follow-up measurement (≤3 months post initiation of intervention)?	0.5
ii) Long-term follow-up measurement (>3 months post initiation of intervention)?	0.5
J Sample size for each group described at randomisation and all important outcome assessments?	1
K Point estimates and measures of variability presented for primary and secondary outcome measures?	1
L i) Sampling method adequate?	0.5
ii) Participation rate adequate (>80%)?	0.5
DQ score	6
Total score (IV+DQ scores)	12



**Figure 1** Included trials at study selection, pooling, synthesis of evidence and sensitivity analyses.

(positive and negative) are reported in two papers describing the study (table 3). Considered together, there is conflicting evidence for the efficacy of exercise interventions.

### Manual handling interventions

Eight trials of manual handling interventions applied as a singular strategy were identified: two RCTs and six NCTs (table 4). The interventions included manual handling training at the workplace<sup>29 35 38</sup> and during nursing studies,<sup>11 32</sup> and the provision of manual handling equipment and training.<sup>20 31 36</sup> All trials showed no effect with the exception of the RCT by Yassi *et al*<sup>36</sup> which showed a reduction in LBP in both intervention groups, although no change in injury or disability. When the levels of evidence ratings are applied, there is conflicting evidence for provision of manual handling equipment and training, and moderate evidence that manual handling training alone (in the workplace or during nurse training) is not effective in preventing back pain.

### Lumbar supports

A single trial was identified that tested the efficacy of lumbar supports. Allen and Wilder's<sup>34</sup> RCT of back belt utilisation during patient transfers showed a positive effect with less back injury in study nurses (table 5). This provides limited evidence

that lumbar supports are effective in preventing back injury in nurses.

### Stress management

The stress management intervention tested by Horneij and colleagues<sup>27</sup> in a high quality trial did not reduce LBP (table 5). Therefore, there is moderate evidence that stress management interventions are not effective in preventing LBP in nurses.

### Multidimensional interventions

Three trials applied combined or multidimensional intervention approaches (table 5). Alexandre *et al*<sup>37</sup> examined a combined manual handling and exercise intervention and found a reduction in LBP frequency and intensity. Linton and colleagues<sup>33</sup> applied an intensive residential program of exercise, manual handling training, pain and lifestyle management and risk assessment training that resulted in a reduction in LBP intensity. Yassi *et al*<sup>21</sup> implemented an early intervention post injury program that involved management by a physician, physiotherapist and occupational therapist and modified duties on return to work. It showed a positive effect on LBP and back injury. The evidence from these singular trials provides limited evidence that each multidimensional intervention strategy is effective. Considered together, there is moderate evidence that

**Table 2** Methodological quality assessment results

Trials	Internal validity (IV) criteria							IV score	Descriptive quality (DQ) criteria								DQ score	Total score
	A	B	C	D	E(i)	E(ii)	F		G	H	I(i)	I(ii)	J	K	L(i)	L(ii)*		
Alexandre <i>et al</i> <sup>37</sup>	+	?	?	+	-	+	+	3.5/6	+	+	-	+	+	+	-	NA	4.5/6	8.0/12
Allen and Wilder <sup>34</sup>	+	-	+	+	-	+	+	4.5/6	-	-	-	+	+	-	-	NA	1.5/6	6.0/12
Besi <sup>38</sup>	+	+	+	-	-	-	+	4.0/6	+	-	+	+	+	-	+	+	4.0/6	8.0/12
Dehlin <i>et al</i> <sup>28</sup>	-	-	-	-	-	-	-	0/6	+	-	+	-	+	-	-	NA	2.5/6	2.5/12
Dehlin <i>et al</i> <sup>29</sup>	-	-	?	-	-	-	-	0/6	+	+	+	-	+	-	-	NA	3.5/6	3.5/12
Gundewall <i>et al</i> <sup>30</sup>	+	+	?	-	-	+	-	2.5/6	+	+	-	+	+	-	?	?	4.0/6	6.5/12
Hellsing <i>et al</i> <sup>32</sup>	-	-	?	?	-	?	?	0/6	+	+	-	+	-	+	?	?	3.0/6	3.0/12
Horneij <i>et al</i> <sup>27</sup>	+	+	+	-	+	-	+	4.5/6	+	+	-	+	+	+	+	+	5.5/6	10.0/12
Knibbe and Friele <sup>20</sup>	-	-	-	-	-	+	+	1.5/6	+	-	-	+	-	+	+	+	2.5/6	4.0/12
Linton <i>et al</i> <sup>33</sup>	+	+	?	-	-	+	-	2.5/6	+	+	+	+	-	-	NA	3.0/6	5.5/12	
Smedley <i>et al</i> <sup>31</sup>	-	-	?	-	-	-	-	0/6	+	-	-	+	+	+	-	-	4.0/6	4.0/12
Videman <i>et al</i> <sup>31</sup>	-	-	+	?	-	-	+	2.0/6	+	-	-	+	+	-	+	+	3.5/6	5.5/12
Wigaeus Hjelm <i>et al</i> <sup>25</sup>	?	?	?	+	-	+	-	1.5/6	+	-	-	+	+	-	?	?	3.0/6	4.5/12
Wood <sup>35</sup>	-	-	?	-	-	?	-	0/6	+	+	-	+	-	+	+	+	3.5/6	3.5/12
Yassi <i>et al</i> <sup>21</sup>	-	-	-	+	-	+	-	1.5/6	+	+	-	+	+	-	+	-	4.0/6	5.5/12
Yassi <i>et al</i> <sup>36</sup>	?	?	?	?	-	+	+	1.5/6	+	+	-	+	+	-	+	+	4.5/6	6.0/12

+, yes; -, no; ?, don't know.

\*If score "no" for L(i), then L(ii) is NA.

multidimensional interventions are effective in preventing LBP in nurses.

### Secondary outcomes

Secondary outcomes such as functional status and sick leave were less frequently measured. There were no cases of multiple trials for any similar intervention and outcome. Gundewall and colleagues<sup>30</sup> physiotherapist-led exercise program provided limited evidence for a reduction in time loss from work. The home exercise program of Horneij *et al*<sup>27</sup> provided moderate evidence for a reduction in LBP interference with work and/or leisure activities, whilst the stress management program tested in the same study was ineffective in this regard. The combined exercise and behavioural therapy intervention of Linton and colleagues<sup>33</sup> provided limited evidence for an increase in satisfaction with activities of daily living. Finally, the early intervention program by Yassi *et al*<sup>21</sup> provided limited evidence for reduction in disability and time loss from work.

### Sensitivity analyses

Figure 1 shows the number of trials included during the sensitivity analyses. During the first sensitivity analysis, the cut-off for high quality status was reduced (see Methods), resulting in two further trials being classified as high quality. These include the multidimensional manual handling and exercise intervention of Alexandre *et al*<sup>37</sup> and the manual handling training intervention by Best.<sup>38</sup> There were no changes to overall evidence findings as a result of these changes. In the second sensitivity analysis, nine trials were excluded due to scoring less than 2/6 on internal validity criteria<sup>20 21 25 28 29 31 32 35 36</sup> leaving seven eligible trials. There was one change to the evidence grading. We found moderate evidence that the provision of manual handling equipment and education is not effective (compared with conflicting evidence previously). The minor changes in evidence findings during the

sensitivity analyses indicate the results of this review are robust and not greatly affected by alterations in review methodology.

## DISCUSSION

This review was needed to synthesise current evidence in relation to back pain and injury prevention in nurses. A lack of high quality studies and infrequent trial replication resulted in no strong evidence for or against any intervention method. Whilst no definitive statements can be made, moderate evidence from multiple trials suggests that multidimensional strategies are effective and manual handling training in isolation is ineffective. For all other interventions there is conflicting evidence or only single trials are available. The specific findings of the review merit discussion in relation to current policy and evidence from other populations.

### Intervention efficacy

Our review concluded that there is moderate evidence that manual handling training alone does not prevent back pain in nurses. This finding is consistent with a review which includes data from multiple worker populations.<sup>39</sup> Hignett<sup>16</sup> found strong evidence that manual handling interventions based predominantly on technique training do not reduce musculoskeletal injury rates. Many nursing workplaces provide annual manual handling training to minimise LBP and prevent workplace back injury. The evidence from this review and that performed by others<sup>16 39</sup> indicates that workplace policies that involve this strategy in isolation are not supported by evidence and should be reconsidered.

Back pain is complex and multifaceted, and the application of multidimensional interventions has been recommended by others.<sup>15</sup> We found consistent positive findings to support the efficacy of multidimensional strategies in preventing LBP in nurses.<sup>21 33 37</sup> Although interventions varied greatly in design and content, when considered together there was moderate evidence supporting multidimensional programs. A synthesis of

**Table 3** Trials of exercise interventions

Study (IV score)	Methods	Subjects	Intervention	Outcomes	Results
<i>Exercise at work</i>					
Gundewall <i>et al</i> <sup>30</sup> (2.5/6)	RCT: stratified then random allocation to 2 groups. Follow-up 13 months	69 nurses/nursing aides with and without LBP working in hospital	(1) 20-min strength, endurance, coordination exercise (6 sessions/month for 13 months); (2) no intervention	LBP intensity, number of days with LBP, lost work days due to LBP	Reduction of LBP prevalence ( $p < 0.02$ ), LBP intensity ( $p < 0.04$ ) and lost work days ( $p < 0.01$ )
Wigaeus Hjelm <i>et al</i> , <sup>25</sup> Josephson <i>et al</i> <sup>26</sup> (1.5/6)	RCT: random allocation to 3 groups. Follow-up 6 months	131 nursing aides working in hospital	All sessions 40 min, 2×/week for 6 months: (1) bicycle ergometer; (2) 3×15 repetitions on 7 equipments; (3) education in occupational health and stress management	LBP intensity	Unclear: Josephson <i>et al</i> <sup>26</sup> report no significant differences; Wigaeus Hjelm <i>et al</i> <sup>25</sup> report reduced LBP in (1) and (2) vs (3) in those with LBP at baseline
Dehlin <i>et al</i> <sup>28</sup> (0/6)	NCT: allocation by work building to 3 groups. Follow-up 8 weeks	46 female nursing aides with LBP working in hospital	2×/week for 8 weeks: (1) strengthening exercise; (2) 30-min lectures on medicine and nursing care; (3) no intervention	Frequency, intensity, duration of LBP and influence of LBP on working capacity	Reduction in LBP duration in (1) vs (2). No other significant differences
Dehlin <i>et al</i> <sup>29</sup> (0/6)	NCT: allocation by work building to 3 groups. Follow-up 8 weeks	45 female nursing aides with LBP working in hospital	2×/week for 8 weeks: (1) 45-min endurance and aerobic exercise; (2) manual handling training; (3) no intervention	Frequency, intensity, duration of LBP and influence of LBP on working capacity	No significant differences
<i>Exercise at home</i>					
Horneij <i>et al</i> <sup>27</sup> (4.5/6)	RCT: random allocation to 3 groups. Follow-up 12 and 18 months	282 female nursing aides working in home-care services	(1) Posture, balance, endurance, functional, stretching and cardiovascular exercises (suggested $\geq 2 \times$ /week); (2) stress reduction training 7×1.5 h plus follow-up; (3) "live as usual"	LBP prevalence, interference with activities, pain drawing	No significant differences in LBP, but (1) had less activity interference compared with (3) at 12-month follow-up

IV, internal validity.

systematic review evidence by Burton and colleagues<sup>39</sup> supported the efficacy of multidimensional interventions for preventing LBP in worker populations, although it was not suggested what the content of these interventions should be and effect size was thought to be potentially modest.

Based on the literature assessed in this study, there is conflicting evidence for interventions that involve the provision of manual handling equipment and education. The RCT by Yassi *et al*<sup>36</sup> included a “no strenuous lifting” group (group 1) and a “safe lifting” group (group 2) and found a reduction in LBP in both groups. The other trials showed no effect,<sup>20–31</sup> although notably the trial by Smedley and colleagues<sup>31</sup> had marked methodological flaws.

In the last decade, nursing associations have developed and advocated “no lift” policies, where equipment is provided and manual lifting is eliminated except in exceptional circumstances.<sup>40</sup> A recent evaluation of a “no lift” program in Australian hospitals reported “strong evidence” that it reduced back injury, pain symptoms and sick leave in nursing staff.<sup>40</sup> This claim is questionable, since the evaluation was cross-sectional and differences between groups such as full-time work and years in the profession (an independent risk factor for back pain)<sup>5</sup> could have influenced findings. An evaluation of a multifaceted ergonomics program that included a “no lift” policy in the United States also reported positive findings<sup>41</sup> but was evaluated using a non-controlled pre-post design that is

inherently susceptible to bias. Of the three trials included in this review, the “no strenuous lifting” intervention of Yassi and colleagues<sup>36</sup> was most closely aligned to the “no lift” strategy and reported a positive effect in relation to LBP although no change in injury or disability outcomes. The other trials provided equipment and education but did not specifically stipulate that manual lifting should be eliminated.<sup>20–31</sup> Hence the popular “no lift” policy may have some efficacy in nursing populations, but there is no consistent evidence presently available to support its widespread application.

We found conflicting evidence for the efficacy of exercise to prevent LBP in nurses. Two trials were highly susceptible to bias, with scores of 0/6 for internal validity.<sup>28–29</sup> The results for one RCT were unclear with conflicting findings reported in two papers describing the same study.<sup>25–26</sup> In the sensitivity analysis, lowest quality evidence was omitted leading to consideration of two trials: an individualised home exercise program did not reduce LBP prevalence,<sup>27</sup> and physiotherapist-led exercise in the workplace resulted in reduced LBP prevalence and intensity.<sup>30</sup> The findings suggest that self-paced exercise programs performed in nurses’ leisure time may not be as effective as directed and physiotherapist-led programs undertaken in the workplace. The ineffectiveness of the exercise program of Horneij and colleagues<sup>27</sup> could possibly be due to poor compliance, since the exercise group did not perform home exercises more frequently than comparison groups. The

**Table 4** Trials of manual handling interventions

Study (IV score)	Methods	Subjects	Intervention	Outcomes	Results
<i>Equipment provision and training</i>					
Yassi <i>et al</i> <sup>36</sup> (1.5/6)	RCT: random allocation to 3 groups. Follow-up 6 and 12 months	346 nurses/nursing aides working in hospital	(1) Sliding and transfer equipment, one lifter, 3-h MH training; (2) multiple new lifters, sit-stand lifters and sliding devices, 3-h MH training; (3) usual practice	Work-related LBP, disability, back injuries	Compared with (3), LBP reduced in (2) at 6 months ( $p=0.02$ ) and in (1) at 12 months ( $p=0.04$ ). No change in injury or disability
Knibbe and Friele <sup>20</sup> (1.5/6)	NCT: incomplete random allocation of 20 teams to 2 groups. Follow-up 12 months	355 female nurses working in home care services	(1) 40 hoists, 12 lifting coordinators, 4-h MH training; (2) usual practice (2 hoists)	Back pain prevalence	No significant differences
Smedley <i>et al</i> <sup>31</sup> (0/6)	NCT: allocation by hospital to 2 groups. Follow-up 32 months	Female nurses working in hospital: 1157 at baseline and 1078 at follow-up	(1) 700 sliding sheets, additional MH equipment and training; (2) no study intervention, but hospital improved MH training and equipment during study	Prevalence of LBP lasting more than a day during the past month	No significant differences
<i>Training in the workplace</i>					
Best <sup>38</sup> (4/6)	RCT: random allocation by nursing home to 3 groups. Follow-up 3 and 12 months	55 female nurses working in nursing homes	(1) 32-h MH training; (2) and (3): in-house orientation training only	Back pain frequency and severity, back injury	No significant differences
Dehlin <i>et al</i> <sup>29</sup> (0/6)	NCT: allocation by work building to 3 groups. Follow-up 8 weeks	45 female nursing aides with LBP working in hospital	2×/week for 8 weeks: (1) 45-min endurance and aerobic exercise; (2) MH training; (3) no intervention	Frequency, intensity, duration of LBP and influence of LBP on working capacity	No significant differences
Wood <sup>35</sup> (0/6)	NCT: allocation by hospital unit to 2 groups. Follow-up 12 months	42 registered nurses and 135 nursing aides working in hospital	(1) 30-min task observation and feedback, 1-h MH training; (2) no intervention	Back injury	No significant differences
<i>Training during nursing studies</i>					
Videman <i>et al</i> <sup>11</sup> (2/6)	NCT: allocation by year of enrolment to 2 groups. Follow-up 12 months	308 student nurses	(1) 40-h training in biomechanics and ergonomics over 2.5 years; (2) usual curriculum	Back pain incidence, severity, injury and disability	No significant differences
Hellsing <i>et al</i> <sup>22</sup> (0/6)	NCT: allocation by school to 2 groups. Follow-up 12, 24 and 36 months	52 student nurses	(1) 2 h/week ergonomics education over 2 years; (2) usual curriculum (5 h ergonomics education)	Annual LBP	No significant differences

IV, internal validity; MH, manual handling.

conflicting findings in the two studies could also be related to differences in the frequency and type of exercises performed.

European guidelines report high level evidence to recommend physical exercise as an intervention for prevention of LBP in general worker populations.<sup>39</sup> Other systematic reviews report limited<sup>43 44</sup> and strong<sup>45 46</sup> evidence to support the use of exercise interventions in preventing LBP, with a meta-analysis indicating a moderate effect size (0.53).<sup>43</sup> The type of exercise that is most effective is not known. One review recommended strengthening and stretching exercises over other strategies,<sup>47</sup> and the successful interventions in another review commonly utilised strengthening and trunk stabilising exercises.<sup>46</sup> Programs of specific stabilisation exercises involving deep spinal and abdominal muscles have recently been shown to be more effective than usual medical care and education in treating chronic LBP, and as effective as spinal manipulative therapy.<sup>48</sup>

One high quality trial in this review tested the efficacy of a stress management program and provided moderate evidence that this intervention in isolation is ineffective in preventing LBP in nurses.<sup>27</sup> It could be argued that stress is not an important risk factor for back pain, as there is inconsistent evidence for an association between job stress and back disorders in nursing cohorts.<sup>5</sup> Systematic reviews of the role of occupational stress in back pain in general populations are also inconsistent: Linton<sup>49</sup> found strong evidence that stress is related to future back pain, but Hartvigsen *et al*<sup>50</sup> disputed this conclusion based on their finding of insufficient evidence. A lack of trial replication inhibits the conclusions that can be drawn regarding the effectiveness of stress management in preventing back pain and injury in nurses.

We found limited evidence from a single trial to support the use of lumbar supports to prevent back injury in nurses. This is in contrast to other reviews that found lumbar supports are ineffective in preventing occupational LBP<sup>43 45 51</sup> or found inconsistent results.<sup>52</sup> European guidelines state that back

belts/lumbar supports are not recommended for prevention of LBP in workers.<sup>39</sup> Whether nurses are unique in their response to lumbar support during patient handling tasks is unknown due to a lack of multiple trials.

### Limitations

There are a number of potential limitations in this review that warrant consideration. Firstly, whilst the literature search was comprehensive and included 10 relevant databases, theses and manual searching of reference lists, the slim possibility that published trials were missed should be acknowledged. Publication bias relating to publishing positive findings may also have played a role. Bias due to publication language was potentially present with one trial ineligible due to reporting in Finnish.<sup>19</sup> It is not known whether the trial, which appeared relevant based on abstract information available in the English language, would have met selection criteria following review of the full paper. Notably, the trial evaluated the efficacy of an exercise intervention, and since this review found conflicting evidence in relation to exercise, additional data from the Finnish study (or the lack thereof) are unlikely to have influenced the conclusions of this review.

Further possible bias in systematic reviews relates to the provision of adequate information by authors of papers to enable grading of methodological quality. In some cases authors of trials included in this review were not able to provide information to the review team (such as when data were no longer accessible). In other cases authors did not respond to requests for information. This may have impacted upon methodological quality scores and the review findings.

It was not possible to present quantitative data on the magnitude of effect of examined interventions in this review. Meta-analyses of trials in each intervention subgroup were considered inappropriate by the reviewing team due to the methodological heterogeneity and poor quality of included studies. Data pooling from low quality studies could compound

**Table 5** Trials of lumbar supports, stress management and multidimensional interventions

Study (IV score)	Methods	Subjects	Intervention	Outcomes	Results
<i>Lumbar supports</i>					
Allen and Wilder <sup>34</sup> (4.5/6)	RCT: random allocation to 2 groups. Follow-up 6 months	50 licensed practical nurses working in hospital	(1) Training in back belts, wearing belts when lifting, 3-h MH training; (2) 3-h MH training only	Incident back injuries and total hours lost due to injury	Reduced incidence of back injuries, no analysis of lost work hours reported
<i>Stress management</i>					
Horneij <i>et al</i> <sup>27</sup> (4.5/6)	RCT: random allocation to 3 groups. Follow-up 12 and 18 months	282 female nursing aides working in home care services	(1) Home exercise program; (2) stress reduction training of 7×1.5 h plus 3- and 6-month follow-up; (3) "live as usual"	LBP prevalence, interference with activities, pain drawing	No significant differences for stress management program
<i>Multidimensional interventions</i>					
Alexandre <i>et al</i> <sup>37</sup> (3.5/6)	RCT: random allocation to 2 groups. Follow-up 4 months	56 female nursing aides with back pain working in hospital	(1) Strength and flexibility exercise, relaxation and MH education (1 h, 2×/week for 4 months); (2) 45-min lecture on anatomy and MH	LBP frequency and intensity (7-day and 2-month)	Reduced LBP frequency (7-day) (p=0.07) and intensity (7-day and 2-month)
Linton <i>et al</i> <sup>33</sup> (2.5/6)	RCT: random allocation to 2 groups. Follow-up post-intervention and 6 months	66 female licensed practical nurses/nursing aides working in hospital who had taken sick leave for back pain in previous 2 years	(1) 5×40 h/week residential program (4 h exercise/day, MH training, pain and lifestyle management and risk assessment); (2) waiting list	Daily LBP intensity, satisfaction with ADL, days lost due to sick leave	Reduced LBP intensity (p=0.01), greater satisfaction with ADL post-intervention (p<0.01), no significant change in sick leave
Yassi <i>et al</i> <sup>21</sup> (1.5/6)	NCT: allocation to 2 groups based on ward. Follow-up 6 months	183 registered nurses working in hospital who sustained an acute back injury during the study	(1) Post injury program: physician, physiotherapy, occupational therapy if >4 days off work, modified duties; (2) routine care	LBP, disability, back injuries and lost work time	Reduced LBP, disability, back injuries, lost-time back injuries and total work time lost (p<0.01)

ADL, activities of daily living; IV, internal validity; MH, manual handling.

any biases present in individual studies and result in an erroneous and misleading pooled estimate. In any case, meta-analyses could not be performed as only three of 16 included studies reported data necessary to calculate effect sizes.

Finally, whilst the database search terms and criteria used to assess methodological quality were based on those recommended by the editorial board of a leading back review group,<sup>17 18</sup> and have been used in other systematic reviews,<sup>53 54</sup> the possibility remains that bias was introduced by the criteria selected for this analysis.

### Implications for research

The overall poor quality of trials undertaken on interventions to prevent back pain and injury in nursing cohorts is concerning given that this is a high risk population with frequent and expensive work absenteeism and workers' compensation.<sup>2 13 14</sup> Although many intervention studies are reported in the literature, a large number of trials were ineligible for this review due to inadequacies of the control group. Further, most eligible trials scored poorly on internal validity criteria, indicating a high susceptibility to bias.

Health economic evaluation of preventative programs has infrequently been undertaken, with the exception of two trials: Yassi *et al*<sup>21</sup> found that combined medical and compensation expenditure was significantly reduced in back-injured nurses who underwent a multidisciplinary therapy program, and Gundewall *et al*<sup>30</sup> found a cost-benefit ratio greater than 10 in their trial of physiotherapist-led exercise. Cost-benefit analyses that contrast intervention expenses with workplace efficiency gains and savings from reduced workers' compensation and work absenteeism are important to inform decisions relating to policy.

Adequately powered high quality randomised controlled studies are needed to answer important research questions about the efficacy of interventions to prevent back pain and back injury in nurses. Internal validity must be maximised, outcome instruments must have proven reliability and validity, and reporting must be comprehensive. Outcome measures should be consistent and include a range of measures such as back pain frequency and severity, injury, functional status, time loss from work and cost-benefit ratio.

Multidimensional interventions show consistent efficacy and warrant further investigation. The most appropriate components are not known, and different composite interventions should be tested with comparison groups. Manual handling training as a singular strategy has not been effective and appears unworthy of further examination. There is conflicting evidence for the efficacy of exercise. Consistent positive findings in other populations suggest that it is a strategy worthy of further study, and examination of different forms of exercise are needed to identify which, if any, are effective in nurses. Due to conflicting evidence, further research is needed to determine whether manual handling interventions that incorporate equipment and training, such as "no lift" strategies, are effective. Only single trials have been undertaken on the effectiveness of stress management and lumbar supports in nursing cohorts, resulting in limited data available to evaluate their efficacy. Systematic review evidence from other working populations might be used to guide research planning decisions for these intervention methods. Other intervention strategies not previously investigated with nurses might also be considered.

Since nurse qualifications, work settings and job tasks can vary considerably, trials should be replicated with different grades of nurses and in different work environments to investigate consistency of results. Nurses with pre-existing back pain and injury and those with no history of symptoms

should be included in the participant cohort to investigate primary and secondary preventative effects.

### CONCLUSION

It is clear from this review that there is a paucity of quality evidence for interventions that aim to prevent back pain and back injury in nurses. This is disappointing, since work absenteeism and workers' compensation due to back disorders in nursing cohorts are both common and expensive. It is critical that policy makers become aware of the current evidence in relation to preventative interventions: data from multiple trials suggest that multidimensional strategies are effective and manual handling training in isolation is ineffective; however, there is no strong evidence to support any firm conclusions. For all other interventions (exercise, lumbar supports, stress management and manual handling equipment and training) there is conflicting evidence or only single trials have been undertaken. Low quality studies do little to answer research questions and should not be endorsed. This review highlights the need for adequately powered randomised controlled studies to provide high quality evidence regarding the effectiveness of interventions to prevent back pain and injury in nursing populations.

### ACKNOWLEDGEMENTS

We acknowledge WorkCover Corporation of South Australia for partial funding of this review. We thank the authors of studies included in this review for providing information to enable grading of methodological quality.

#### Main messages

- A systematic literature review identified 16 predominantly low quality trials of interventions that aim to prevent back pain or back injury in nursing cohorts.
- There is moderate evidence from multiple trials that manual handling training in isolation is not effective and multidimensional strategies are effective.
- There is conflicting evidence regarding the efficacy of exercise and the provision of manual handling equipment and training as preventative measures.
- Single trials provide limited evidence that lumbar supports are effective and moderate evidence that stress management programs are not effective.
- High quality research is urgently needed to investigate conflicting findings and further examine the efficacy of interventions in preventing back pain and back injury in nurses.

#### Policy implications

- Health service organisations must become aware of the evidence in relation to back pain and injury prevention in nurses and design policy accordingly.
- Evidence from multiple trials suggests that manual handling training in isolation is not effective in preventing back pain and injury in nurses.
- Multidimensional strategies show consistent positive outcomes although the most important intervention components are not known.

**Authors' affiliations**

**Anna P Dawson, Gwendolen A Jull, Paul W Hodges**, Division of Physiotherapy, School of Health and Rehabilitation Sciences, The University of Queensland, Brisbane, Australia

**Skye N McLennan**, School of Psychology, University of Adelaide, Adelaide, Australia

**Stefan D Schiller**, Division of Health Sciences, University of South Australia, Adelaide, Australia

**Simon Stewart**, Baker Heart Research Institute, Melbourne, Australia

Funding: WorkCover Corporation of South Australia provided financial support towards the completion of this systematic review. Professor Paul Hodges is supported by the National Health and Medical Research Council of Australia. The funding bodies had no role in designing or undertaking the review.

Competing interests: None.

**REFERENCES**

- Hofmann F, Stossel U, Michaelis M, et al. Low back pain and lumbago-sciatica in nurses and a reference group of clerks: results of a comparative prevalence study in Germany. *Int Arch Occup Environ Health* 2002;**75**:484-90.
- Cohen-Mansfield J, Culpepper WJ, Carter P. Nursing staff back injuries: prevalence and costs in long term care facilities. *AAOHN J* 1996;**44**:9-17.
- Heliövaara M. Occupation and risk of herniated lumbar intervertebral disc or sciatica leading to hospitalization. *J Chronic Dis* 1987;**40**:259-64.
- Bureau of Labor Statistics. Workplace injuries and illnesses in 2005. Washington, DC: United States Department of Labor, 2006, <http://www.bls.gov/news.release/pdf/osh.pdf> (accessed 25 June 2007).
- Sherehiy B, Karwowski W, Marek T. Relationship between risk factors and musculoskeletal disorders in the nursing profession: a systematic review. *Occupational Ergonomics* 2004;**4**:241-79.
- Baldasseroni A, Tartaglia R, Sgarrella C, et al. Frequenza della lombalgia in una coorte di allievi infermieri. *Med Lav* 1998;**89**:242-53.
- Ready AE, Borenskie SL, Law SA, et al. Fitness and lifestyle parameters fail to predict back injuries in nurses. *Can J Appl Physiol* 1993;**18**:80-90.
- Feyer A, Herbison P, Williamson AM, et al. The role of physical and psychological factors in occupational low back pain: a prospective cohort study. *Occup Environ Med* 2000;**57**:116-20.
- Mostardi R, Noe D, Kovacic M, et al. Isokinetic lifting strength and occupational injury. A prospective study. *Spine* 1992;**17**:189-93.
- Klüber Moffett JA, Hughes GI, Griffiths P. A longitudinal study of low back pain in student nurses. *Int J Nurs Stud* 1993;**30**:197-212.
- Videman T, Rauhala H, Asp S, et al. Patient-handling skill, back injuries and back pain. An intervention study in nursing. *Spine* 1989;**14**:148-56.
- Adams M, Mannion A, Dolan P. Personal risk factors for first-time low back pain. *Spine* 1999;**24**:2497-505.
- Bureau of Labor Statistics. Case and demographic characteristics for work-related injuries and illnesses involving days away from work. Resource table 10: detailed occupation by selected parts of body affected. United States Department of Labor, 2002. <http://www.bls.gov/iif/oshwc/osh/case/ostbl277.pdf> (accessed 27 June 2007).
- WorkCover Corporation of South Australia. *Nurses' back injury claims 1999-2002*. Adelaide, South Australia: WorkCover Corporation of South Australia, 2004.
- Lagerstrom M, Hansson T, Hagberg M. Work-related low-back problems in nursing. *Scand J Work Environ Health* 1998;**24**:449-64.
- Hignett S. Intervention strategies to reduce musculoskeletal injuries associated with handling patients: a systematic review. *Occup Environ Med* 2003;**60**:e6.
- van Tulder M, Assendelft WJJ, Koes B, et al. Method guidelines for systematic reviews in the Cochrane Collaboration Back Review Group for Spinal Disorders. *Spine* 1997;**22**:2323-30.
- van Tulder M, Furlan A, Bombardier C, et al. Updated method guidelines for systematic reviews in the Cochrane Collaboration Back Review Group. *Spine* 2003;**28**:1290-9.
- Karvonen MJ, Jarvinen T, Nummi J. Seurantatutkimus sairaanhoitajien selkävaikeista follow-up study on the back problems of nurses. Helsinki, Finland: Institute of Occupational Health, 1977.
- Knibbe JJ, Friele RD. The use of logs to assess exposure to manual handling of patients, illustrated in an intervention study in home care nursing. *International Journal of Industrial Ergonomics* 1999;**24**:445-54.
- Yassi A, Tate R, Cooper JE, et al. Early intervention for back-injured nurses at a large Canadian tertiary care hospital: an evaluation of the effectiveness and cost benefits of a two-year pilot project. *Occup Med* 1995;**45**:209-14.
- Cooper JE, Tate RB, Yassi A, et al. Effect of an early intervention program on the relationship between subjective pain and disability measures in nurses with low back injury. *Spine* 1996;**21**:2329-36.
- Cooper JE, Tate RB, Yassi A. Components of initial and residual disability after back injury in nurses. *Spine* 1998;**23**:2118-22.
- Tate RB, Yassi A, Cooper J. Predictors of time loss after back injury in nurses. *Spine* 1999;**24**:1930-5.
- Wigaeus Hjelm E, Hagberg M, Hellstrom S. Prevention of musculoskeletal disorders in nursing aides by physical training. In: Hagberg M, Hofmann F, Stossel U, et al, eds. *Occupational health for health care workers*. Freiburg, Germany: Ecomed, 1992:364-6.
- Josephson M, Hagberg M, Hjelm EW. Self-reported physical exertion in geriatric care. A risk indicator for low back symptoms? *Spine* 1996;**21**:2781-5.
- Horneij E, Hemborg B, Jensen I, et al. No significant differences between intervention programmes on neck, shoulder and low back pain: a prospective randomized study among home-care personnel. *J Rehabil Med* 2001;**33**:170-6.
- Dehlin O, Berg S, Hedenrud B, et al. Muscle training, psychological perception of work and low-back symptoms in nursing aides. *Scand J Rehabil Med* 1978;**10**:201-9.
- Dehlin O, Berg S, Andersson GB, et al. Effect of physical training and ergonomic counseling on the psychological perception of work and on the subjective assessment of low-back insufficiency. *Scand J Rehabil Med* 1981;**13**:1-9.
- Gundewall B, Liljeqvist M, Hansson T. Primary prevention of back symptoms and absence from work. A prospective randomized study among hospital employees. *Spine* 1993;**18**:587-94.
- Smedley J, Trevelyan F, Inskip H, et al. Impact of ergonomic intervention on back pain among nurses. *Scand J Work Environ Health* 2003;**29**:117-23.
- Hellsing AL, Linton SJ, Andershed B, et al. Ergonomic education for nursing students. *Int J Nurs Stud* 1993;**30**:499-510.
- Linton S, Bradley LA, Jensen I, et al. The secondary prevention of low back pain: a controlled study with follow-up. *Pain* 1989;**36**:197-207.
- Allen S, Wilder K. Back belts pay off for nurses. *Occup Health Saf* 1996;**65**:59-62.
- Wood D. Design and evaluation of a back injury prevention program within a geriatric hospital. *Spine* 1987;**12**:77-82.
- Yassi A, Cooper JE, Tate RB, et al. A randomized controlled trial to prevent patient lift and transfer injuries of health care workers. *Spine* 2001;**26**:1739-46.
- Alexandre N, de Moraes M, Correa Filho H, et al. Evaluation of a program to reduce back pain in nursing personnel. *Rev Saude Publica* 2001;**35**:356-61.
- Best M. An evaluation of manutention training in preventing back strain and resultant injuries in nurses. *Saf Sci* 1997;**25**:207-22.
- Burton A, Balague F, Cardon G, et al. How to prevent low back pain. *Best Pract Res Clin Rheumatol* 2005;**19**:541-55.
- Engkvist I. Evaluation of an intervention comprising a no lifting policy in Australian hospitals. *Appl Ergon* 2006;**37**:141-8.
- Nelson A, Matz M, Chen F, et al. Development and evaluation of a multifaceted ergonomics program to prevent injuries associated with patient handling tasks. *Int J Nurs Stud* 2006;**43**:717-33.
- Josephson M, Lagerstrom M, Hagberg M, et al. Musculoskeletal symptoms and job strain among nursing personnel: a study over a three year period. *Occup Environ Med* 1997;**54**:681-5.
- van Poppel M, Hooftman W, Koes B. An update of a systematic review of controlled clinical trials on the primary prevention of back pain at the workplace. *Occup Med (Lond)* 2004;**54**:345-52.
- Maher C. A systematic review of workplace interventions to prevent low back pain. *Aust J Physiother* 2000;**46**:259-69.
- Linton S, van Tulder M. Preventive interventions for back and neck pain problems: what is the evidence? *Spine* 2001;**26**:778-87.
- Hayden J, van Tulder M, Malmivaara A, et al. Exercise therapy for treatment of non-specific low back pain. *Cochrane Database Syst Rev* 2005;(3):CD000335.
- Hayden J, van Tulder M, Tomlinson G. Systematic review: strategies for using exercise therapy to improve outcomes in chronic low back pain. *Ann Intern Med* 2005;**142**:776-85.
- Ferreira P, Ferreira M, Maher C, et al. Specific stabilisation exercise for spinal and pelvic pain: a systematic review. *Aust J Physiother* 2006;**52**:79-88.
- Linton S. Occupational psychological factors increase the risk for back pain; a systematic review. *J Occup Rehabil* 2001;**11**:53-66.
- Hartvigsen J, Lings S, Leboeuf-Yde C, et al. Psychosocial factors at work in relation to low back pain and consequences of low back pain: a systematic, critical review of prospective cohort studies. *Occup Environ Med* 2004;**61**:e2.
- van Tulder M, Jellema P, van Poppel M, et al. Lumbar supports for prevention and treatment of low-back pain. *Cochrane Database Syst Rev* 2000;(3):CD001823.
- Ammendolia C, Kerr M, Bombardier C. Back belt use for prevention of occupational low back pain: a systematic review. *J Manipulative Physiol Ther* 2005;**28**:128-34.
- Heymans M, van Tulder M, Esmail R, et al. Back schools for non-specific low-back pain. *Cochrane Database Syst Rev* 2004;(4):CD000261.
- Ostelo R, van Tulder M, Vlaeyen J, et al. Behavioural treatment for chronic low-back pain. *Cochrane Database Syst Rev* 2005;(1):CD002014.